

## Recurrence Relations

**Reading:** Gossett Sections 7.1 and 7.2.

**Definition:** A one-way infinite sequence is a function from the natural numbers to some other set.

E.g.

$$fib(0) = 1, fib(1) = 1, fib(2) = 2, fib(3) = 3, fib(4) = 5, fib(5) = 8, .$$

$$q(0) = 1, q(1) = 2, q(2) = 4, q(3) = 8, \dots$$

$$pr(0) = 2, pr(1) = 3, pr(2) = 5, pr(3) = 7, \dots$$

**Definition:** A recurrence relation is an equation that defines all members of a sequence past a certain point in terms of earlier members. That is an equation

$$a(n) = F, \text{ for all } n \in \{n_1, n_1 + 1, \dots\}$$

where  $F$  is an expression combining only  $a(n - 1)$ ,  $a(n - 2)$ , ...,  $a(0)$ .

**Examples**

- $fib(n) = fib(n - 1) + fib(n - 2)$ , for all  $n \geq 2$
- $q(n) = 2 \times q(n - 1)$ , for all  $n \geq 1$

If we conjoin enough ( $n_1$ ) base cases with the recurrence relation, then together they define a sequence.

**Examples:**

- $fib(0) = 1$
- $fib(1) = 1$
- $q(0) = 1$

## Substitute and simplify method

Consider the sequence defined by

$$\begin{aligned} a(0) &= 5 \\ a(n) &= 2a(n-1) - 3, \text{ for } n \geq 1 \end{aligned}$$

We can reason (rather informally) that.

$$\begin{aligned} &a(n) \\ &= 2a(n-1) - 3 \\ &= 2(2a(n-2) - 3) - 3 \text{ substitution} \\ &= 4a(n-2) - 2 \cdot 3 - 3 \text{ simplify} \\ &= 4(2a(n-3) - 3) - 2 \cdot 3 - 3 \text{ substitution} \\ &= 8a(n-3) - 4 \cdot 3 - 2 \cdot 3 - 3 \text{ simplify} \\ &\quad \vdots \\ &= 2^k a(n-k) - 3(2^{k-1} + 2^{k-2} + \dots + 1) \\ &\quad \vdots \\ &= 2^n a(0) - 3(2^{n-1} + 2^{n-2} + \dots + 1) \\ &= 2^n \cdot 5 - 3 \cdot (2^n - 1) \text{ since } \sum_{i=0}^{n-1} 2^i = 2^n - 1 \\ &= 2 \cdot 2^n + 3 \\ &= 2^{n+1} + 3 \end{aligned}$$

If there is any doubt, we can prove the result by induction. Unfortunately the substitute and simplify method does not always give an easy to simplify result.

## Linear Homogeneous Recurrence Relations with Constant Coefficients of Degree $k$

**Definition:** A *linear homogeneous recurrence relation with constant coefficients* (LHRRCC) is a recurrence relation whose RHS is a sum of terms each of the form

$$c_b \cdot a(n - j)$$

where  $c \in \mathbb{C}$  and  $j \in \mathbb{N}$  are constants.  $\square$

**Definition:** The *degree* of a LHRRCC is the maximum  $b$  value for which the  $c$  value is not 0. I.e. if we have

$$a(n) = c_1 a(n - 1) + c_2 a(n - 2) + \dots + c_k a(n - k)$$

with  $c_k \neq 0$ , then the degree is  $k$ .

### When the degree is 2


Consider the degree 2 case. The RR is

$$a(n) = c_1 a(n - 1) + c_2 a(n - 2) \quad (*)$$

Suppose there is a solution for the recurrence relation of the form

$$a(n) = \theta r^n, \text{ for all } n \in \mathbb{N}$$

for some  $\theta \neq 0$  and some  $r \neq 0$ .



So  $r$  is a root of the polynomial

$$x^2 - c_1 x - c_2 \quad (**)$$

Conversely, if  $r$  is root of the polynomial

$$x^2 - c_1 x - c_2 \quad (**)$$

then for any  $\theta$  and any  $n \in \mathbb{N}$

$$\theta r^n = c_1 \theta r^{n-1} + c_2 \theta r^{n-2}$$

so

$$a(n) = \theta r^n$$

is a solution to (\*).

The characteristic polynomial of (\*) is (\*\*).

---

We have proved the following theorem:

**Theorem:**

If  $r$  is a root of the characteristic polynomial

$$x^2 - c_1x - c_2$$

then, for any  $\theta \in \mathbb{C}$ , the sequence

$$a(n) \triangleq \theta r^n, \text{ for all } n \in \mathbb{N}.$$

is a solution to the equation

$$a(n) = c_1a(n-1) + c_2a(n-2), \text{ for all } n \geq 2.$$

□

**Example.**

$$a(n) = a(n-1) + 6a(n-2)$$

The characteristic polynomial is

$$x^2 - x - 6$$

With roots 3 and  $-2$ .

One root is  $r = 3$ ; picking  $\theta = 1$ , we get a sequence

$$\begin{aligned} &\langle 3^0, 3^1, 3^2, 3^3, \dots \rangle \\ &= \langle 1, 3, 9, 27, \dots \rangle \end{aligned}$$

So if the base cases are  $a(0) = 1$  and  $a(1) = 3$ , we should pick  $r = 3$  and  $\theta = 1$ .

Trying the other root  $r = -2$  and picking  $\theta = 3$  we get

$$\begin{aligned} &\langle 3 \cdot (-2)^0, 3 \cdot (-2)^1, 3 \cdot (-2)^2, 3 \cdot (-2)^3, \dots \rangle \\ &\langle 3, -6, 12, -24, \dots \rangle \end{aligned}$$

So if the base cases are  $a(0) = 3$  and  $a(1) = -6$ , we should pick  $r = -2$  and  $\theta = 3$ .

---

*Unfortunately* the base cases may not allow such a simple solution.

Consider

$$\begin{aligned} a(0) &= -2 \\ a(1) &= 3 \\ a(n) &= a(n-1) + 6a(n-2) \end{aligned}$$

The roots are  $r_1 = 3$  and  $r_2 = -2$ . Trying  $r_1$  we must find  $\theta$  such that

$$\theta 3^0 = -2 \text{ and } \theta 3^1 = 3$$


Trying  $r_2$  we must find a  $\theta$  such that

$$\theta (-2)^0 = -2 \text{ and } \theta (-2)^1 = 3$$

Neither root gives a solution that agrees with both base cases.

---

## Linear combinations of solutions



Suppose that  $w$  and  $x$  are two solutions. I.e.

$$w(n) = c_1 w(n-1) + c_2 w(n-2), \text{ for all } n \geq 2$$


and

$$x(n) = c_1 x(n-1) + c_2 x(n-2), \text{ for all } n \geq 2$$

Then we can build a sequence  $z$  defined by

$$z(n) \triangleq w(n) + x(n), \text{ for all } n \in \mathbb{N}$$

Now  $z$  too will be a solution as, for all  $n \geq 2$



So given any two solutions,  $x$  and  $w$ , any linear combination of them

$$z(n) \triangleq \theta_1 w(n) + \theta_2 x(n), \text{ for all } n \in \mathbb{N}$$

will also be a solution.

A set that is closed under linear combinations is called a *vector space*.

If the degree is 2, all solutions can be formed as linear combination of just 2 (appropriately chosen) solutions. If there are two different roots, we have two solutions  $r_1^n$  and  $r_2^n$ , which will suffice.

## Using both roots at once

Consider, again, the LHRCC of degree 2.

$$a(n) = c_1 a(n-1) + c_2 a(n-2), \text{ for all } n \geq 2 \quad (*)$$

with characteristic polynomial

$$x^2 - c_1 x - c_2 = 0$$


Let  $r_1$  and  $r_2$  be the roots of this polynomial.

Consider any two constants  $\theta_1$  and  $\theta_2$ .

$$\theta_1 r_1^n + \theta_2 r_2^n \text{ is a solution to } (*)$$

since it is a linear combination of the solutions  $r_1^n$  and  $r_2^n$ .

Here is a direct proof:



This proves the following theorem

**Theorem:** Given the LHRCC of degree 2

$$a(n) = c_1 a(n-1) + c_2 a(n-2), \text{ for all } n > 2 \quad (*)$$

If  $r_1$  and  $r_2$  are the two roots of the characteristic polynomial

$$x^2 - c_1 x - c_2$$

then for any  $\theta_1$  and  $\theta_2$

$$\theta_1 r_1^n + \theta_2 r_2^n$$

is a solution to (\*).

---

Furthermore, if we know  $a(0)$  and  $a(1)$ . Then

$$\theta_1 + \theta_2 = a(0)$$

$$\theta_1 r_1 + \theta_2 r_2 = a(1)$$

With 2 linear equations in 2 unknowns we can solve for  $\theta_1$  and  $\theta_2$ .

This will succeed provided  $r_1 \neq r_2$ .

So, when the roots are distinct, we can find the unique values for  $\theta_1$  and  $\theta_2$  that satisfy the two base cases.

**Example:**

$$a(0) = -2$$

$$a(1) = 3$$

$$a(n) = a(n-1) + 6a(n-2)$$

The characteristic polynomial

$$x^2 - x - 6 = 0$$

has roots  $r_1 = 3$  and  $r_2 = -2$ . So we are looking for a solution of the form

$$\theta_1 3^n + \theta_2 (-2)^n$$

We have

$$\theta_1 + \theta_2 = a(0) = -2$$

$$3\theta_1 - 2\theta_2 = a(1) = 3$$

So

$$3\theta_1 - 2(-2 - \theta_1) = 3$$

$$5\theta_1 + 4 = 3$$

$$\theta_1 = \frac{-1}{5}$$

and

$$\theta_2 = -2 - \theta_1$$

$$\theta_2 = \frac{-9}{5}$$

**Procedure**

- From the RR derive the characteristic polynomial
- Find roots  $r_1$  and  $r_2$  of the characteristic polynomial.
- If  $r_1 \neq r_2$  then look for a solution of the form

$$\theta_1 r_1^n + \theta_2 r_2^n$$

- use the base cases to solve for  $\theta_1$  and  $\theta_2$

**Example:**

$$fib(0) = 1$$

$$fib(1) = 1$$

$$fib(n) = fib(n-1) + fib(n-2)$$

Form the characteristic polynomial

$$x^2 - x - 1 = 0$$

Find roots using the quadratic equation. The roots are

$$\frac{1 + \sqrt{5}}{2} = 1.61803 \dots = \phi$$

and

$$\frac{1 - \sqrt{5}}{2} = -0.61803 \dots = 1 - \phi = \frac{-1}{\phi}$$

We are looking for a solution of the form

$$\theta_1 \phi^n + \theta_2 (1 - \phi)^n$$

Using  $fib(0) = fib(1) = 1$  we get

$$\theta_1 + \theta_2 = 1$$

$$\theta_1 \phi + \theta_2 (1 - \phi) = 1$$

Substitute  $\theta_2 = 1 - \theta_1$  into  $\theta_1 \phi + \theta_2 (1 - \phi) = 1$  to get

$$\theta_1 \phi + (1 - \theta_1)(1 - \phi) = 1$$

Now solve for  $\theta_1$ .

$$\theta_1 \phi + (1 - \theta_1)(1 - \phi) = 1$$

$$\theta_1 \phi + (1 - \phi) - \theta_1(1 - \phi) = 1$$

$$\theta_1(\phi - (1 - \phi)) = \phi$$

$$\theta_1(2\phi - 1) = \phi$$

$$\theta_1 \sqrt{5} = \phi$$

$$\theta_1 = \frac{\phi}{\sqrt{5}} = \frac{1 + \sqrt{5}}{2\sqrt{5}} = \frac{5 + \sqrt{5}}{10}$$

And then solve for  $\theta_2$

$$\begin{aligned} \theta_2 &= 1 - \theta_1 \\ &= \frac{\sqrt{5}}{\sqrt{5}} - \frac{\phi}{\sqrt{5}} \\ &= \frac{\sqrt{5} - \phi}{\sqrt{5}} \\ &= \frac{\frac{2\sqrt{5}}{2} - \frac{1 + \sqrt{5}}{2}}{\sqrt{5}} \\ &= \frac{\sqrt{5} - 1}{2} \\ &= \frac{1 - \phi}{\sqrt{5}} \end{aligned}$$

So the solution is

$$\begin{aligned} fib(n) &= \frac{\phi}{\sqrt{5}} \phi^n - \frac{(1 - \phi)}{\sqrt{5}} (1 - \phi)^n \\ &= \frac{1}{\sqrt{5}} \phi^{n+1} - \frac{1}{\sqrt{5}} (1 - \phi)^{n+1} \end{aligned}$$

□

These ideas generalize to degrees larger than 2.



## Repeated roots for LHRCCs with degree 2

When  $r_1 = r_2$  then  $\theta_1 r_1^n + \theta_2 r_2^n$  can be written as  $\theta r^n$  where  $\theta = \theta_1 + \theta_2$  and  $r = r_1 = r_2$ .

So the 2 base cases may form an overdetermined system: 2 equations and one unknown.

### Example

$$a(0) = 1$$

$$a(1) = 2$$

$$a(n) = 6a(n-1) - 9a(n-2), \text{ for } n \geq 2$$

The characteristic polynomial is

$$x^2 - 6x + 9$$

with root  $r = 3$ .

We look for a solution of the form

$$\theta r^n$$

But

$$\theta = 1$$

$$\theta r = 2$$

is not solvable!□

Consider a quadratic with a repeated root

$$x^2 - 4x + 4$$

so  $r = 2$ . This is characteristic of the LHRCC

$$a(n) = 4a(n-1) - 4a(n-2)$$

adding a couple of base cases  $a(0) = 0$  and  $a(1) = 2$  we get

$n$	$a(n)$
0	0
1	2
2	$4 \times 2 = 8$
3	$4 \times 8 - 4 \times 2 = 24$
4	$4 \times 24 - 4 \times 8 = 64$
5	$4 \times 64 - 4 \times 24 = 160$


The solution is apparently  $n2^n$ .

This suggests  $nr^n$  is a potential solution in general.

Note that  $nr^n$  is  $r \times nr^{n-1}$  and that  $nr^{n-1}$  is the derivative of our basic solution  $r^n$ .

So we might do well to look at derivatives.

When a polynomial has a repeated root, that root will also be a root of its derivative:



Consider  $x^2 - 6x + 9 = (x - 3)^2$  and its derivative  $2(x - 3)$ ,  
3 is a root of both.

---

*In general* (for all  $r \in \mathbb{R}$ ) if

$$p(x) = x^2 - c_1x - c_2 = (x - r)^2$$

then  $r$  will also be a root of  $p'(x) = 2x - c_1 = 2(x - r)$ .

Define  $p_0(x) \triangleq x^{n-2} \cdot p(x)$

$r$  will also be a root of  $p_0$  since

$$p_0(r) = r^{n-2} \cdot p(r) = 0.$$

$r$  will be a root of  $p'_0$  since

$$p'_0(r) = r^{n-2} \cdot p'(r) + (n-2)r^{n-3} \cdot p(r) = 0.$$

**Theorem:** If a LHRCC of the form


$$a(n) = c_1a(n-1) + c_2a(n-1), \text{ for all } n \geq 2 \quad (*)$$

has a characteristic polynomial with one root  $r$  then,

$$nr^n$$

is a solution:

**Proof:**





**Theorem:** If a LHRCC of the form

$$a(n) = c_1 a(n-1) + c_2 a(n-1), \text{ for all } n \geq 2 \quad (*)$$

has a characteristic polynomial with one root  $r$  then, for any  $\alpha_0, \alpha_1 \in \mathbb{R}$ ,  $(\alpha_0 + \alpha_1 n)r^n$  is a solution.

**Proof:**  $(\alpha_0 + \alpha_1 n)r^n$  is just a linear combination of the solutions  $r^n$  and  $nr^n$ .  $\square$

We can use the base cases to compute the  $\alpha_0$  and  $\alpha_1$ .

Back to the earlier example

$$a(0) = 1$$

$$a(1) = 2$$

$$a(n) = 6a(n-1) - 9a(n-2), \text{ for } n \geq 2$$

The root of the characteristic polynomial  $x^2 - 6x + 9$  is 3.

From the theorem, the solution is of the form  $(\alpha_0 + \alpha_1 n) \cdot 3^n$

Now solve

$$\alpha_0 = 1$$

$$(\alpha_0 + \alpha_1) \cdot 3 = 2$$

so

$$\alpha_1 = \frac{2}{3} - 1 = -\frac{1}{3}$$

Check:

$n$	$a(n)$	$(1 - \frac{1}{3}n) \cdot 3^n$
0	1	$(1 - \frac{1}{3} \cdot 0) \cdot 3^0 = 1.0$
1	2	$(1 - \frac{1}{3} \cdot 1) \cdot 3^1 = 2.0$
2	$6 \times 2 - 9 \times 1 = 3$	$(1 - \frac{1}{3} \cdot 2) \cdot 3^2 = 3.0$
3	$6 \times 3 - 9 \times 2 = 0$	$(1 - \frac{1}{3} \cdot 3) \cdot 3^3 = 0$
4	$6 \times 0 - 9 \times 3 = -27$	$(1 - \frac{1}{3} \cdot 4) \cdot 3^4 = -27.0$
5	$6 \times -27 - 9 \times 0 = -162$	$(1 - \frac{1}{3} \cdot 5) \cdot 3^5 = -162.0$

**Procedure for LHRCC of degree 2**

- From the RR derive the characteristic polynomial

- Find roots  $r_1$  and  $r_2$  of the characteristic polynomial.
- If  $r_1 \neq r_2$  then look for a solution of the form

$$\theta_1 r_1^n + \theta_2 r_2^n$$

- \* Use the base cases to solve for  $\theta_1$  and  $\theta_2$ .
- If the sole root is  $r$  then look for a solution of the form

$$(\alpha_0 + \alpha_1 n)r^n$$

- \* Use the base cases to solve for  $\alpha_0$  and  $\alpha_1$ .
- 

One can generalize these theorems and the resulting procedure to LHRCCs with any degree and any number of repeated roots. See Gossett's book.